



AIRS Validation Overview & TDS Support of Validation

Eric Fetzer

AIRS Science Team Meeting
November 2001
Pasadena





Today's Talk

- TLSCF Support of AIRS Validation
 - TLSCF (Team Leader Science Computing Facility): the AIRS JPL processing facility
- Matching AIRS / AMSU / HSB observations to correlative data sets
- First Year Validation Schedules

2





TLSCF Data Processing

- Level 0: Instrument Packets from AIRS/AMSU/HSB
 - Pushed from GSFC DAAC (EOS processing center) and archived
- Level 1A: Instrument Counts
 - Generate and archive
- Level 1B: Calibrated Radiances
 - Generate and archive
- Level 2: Retrieved Geophysical Properties
 - Generate and archive about 10%

Match these to validation data sets (More later)



TLSCF Data System: TDS



- A data storage system that includes:
 - A data catalog DOM (Distributed Object Manager)
 - The DOM Interface
 - GUI and command line driven
 - Reflects directory structure -- masochists can cd
 - Tested recently during AIRS science team exercise
 - Virtual links to or direct copy of files
 - Terabytes of storage
 - 0.5 TB cache
 - 2 TB online for <u>TEST</u> data sets
 - 25 TB near-line. Enough for a year -- we hope!



Matchups: Key Validation Files



- AIRS / AMSU / HSB instrument footprints ("golfballs") matched to a specified time and location
 - Independent of validation data type
 - Within specified space-time windows (e. g. 100 km-3 hrs)
 - Contains all AIRS / AMSU / HSB data Levels 1A, 1B & 2
 - Contains pointers to correlative data sets
- Matchup files are archived in the TDS
 - Access is straightforward, with readers available


Matchups involve two data sets --Correlative data in native format, Matchup files in HDF



Matchup Processing



AIRS TLSCF MATCH-UP FILE CREATION AND PROCESSING



November 2001 EJF

AIRS Validation and TDS



Post-meeting: Complete this table



PI	Brief Title	Data Type	Data Size of Order	In TDS Archive?	# Matches	When	Metadata Available?
Atlas	Geophysical Validation	Assimilation analyses	Really Big	No	N/A	TBD	N/A
Barnes	Validation Over Mauna Loa	Lidar	Megabytes	Yes	100	L+2 Months	TBD
Bennartz	'Validation of [MW] at High Latitudes	Microwave radar	Gigabytes?	Limited set	>>100?	Launch?	Probably
McMillan	BBAERI AIRS Ocean: BAOVE	Radiosondes, AERI, lidar	< Gigabytes	Yes	~100	L+2 Months	TBD
Minnett	Trans-Oceanic Measurements	Radiosondes, AERI spectra	< Gigabytes	Yes	~100	TDB by cruises	TBD
Newchurch	Validating AIRS Ozone Observations	Ozonesondes	Megabytes	Yes	~100	L + 2 Months?	TBD
Schmidlin	Temperature, Water Vapor, Ozone Val Measurements	Radiosondes, rocketsondes	Megabytes	Yes	~100	L + 2 Months	TBD
Voemel	'Balloons in upper troposphere	Radiosondes (good H2O)	Megabytes	Yes	~100	November to May is bad	TBD
Walden	Validation ofAIRS over the Antarctic Plateau:	Radiosondes, AERI spectra, lidar	< Gigabytes	Yes	~100	Austral Summer 2001- 02	TBD
Whiteman	Water Vapor and Cloud w/ Lidars and AERI	Lidar, AERI	< GB?	Yes	~100	L + 2 Months	TBD
Yoe	GPS total water	GPS	< GB	Yes	>>100	Launch	TBD







- See John Gieselman for passwords
 - Email: jsg@airs1.jpl.nasa.gov; Phone: (818) 354-7848
- Browser or secure remote login (ssh2) for accessing TDS
 - JPL computer security trust us only slightly more than universities (foreign national particularly problematical).
- Use AIRS Science Team Web Page for communication
 - See http://airsteam.jpl.nasa.gov





AIRS/AMSU/HSB First Year Schedule Rollout and Validation







- Assess AIRS / AMSU / HSB performance 1 year after launch for handover of processing to GSFC DAAC
- Other validation schedule drivers include
 - Instrument activation and spacecraft maneuver schedule
 - Data assimilation impact assessment by first year
- Conceptually, validation activities phased by a sequence of stages
 - Initial instrument activation and commissioning
 - Alignment and geolocation
 - Calibration activities
 - Initial assessment of performance
 - Basic field validation
 - Initiate observations at ground truth sites
 - Assess AIRS radiometric performance compared to SST
 - Assess clear sky detection methods



Drivers and Phasing




(Phasing continued)

- Cloud-clearing validation
 - From clear ocean scenes assess cloud-cleared radiances.
 - Clear sky ground truth for forward model assessment
- Retrieved T,q validation
 - Step 1: local T,q comparisons at well instrumented sites under well characterized conditions
 - Step 2: global performance assessment from RaObs and model analyses
- Extended validation activities
 - Detailed analyses and evaluations using other data sources to continue after initial assessment at Launch + 1 year
 - Validation data acquired after Launch + 8 months will not have impact on initial assessment to be completed at Launch + 12 months - this is aimed to declare data products ready for scientific community and labeled as at least ScienceQAFlag = "Provisional"
 - Later validation will be factored into Launch + 2 year PGE update and data reprocessing with transition of all data products to ScienceQAFlag = "Validated"



AIRS/AMSU/HSB Early Validation Timeline Characterizing the atmospheric column and surface observed by AIRS/AMSU/HSB






AIRS Validation and TDS



AIRS/AMSU/HSB Early Validation Schedule and First Year PGE Deliveries






AIRS Validation and TDS



AIRS Calibration, and L1B - SST Validation








Cloud-Cleared Radiance Validation






November 2001 EJF

AIRS Validation and TDS



Accuracy of Local T and q



November 2001 EJF


AIRS Validation and TDS



EJF

Global T and q -- 1 K/1 km and 20% Humidity





AIRS Validation and TDS

17



Near Term Goals



- Update schedules to include individual investigators' activities
 - Understand schedule constraints to best utilize resources
- Understand correlative data sets to create matchups with AIRS/AMSU/HSB
 - Format, content, readers, metadata...
- Provide further support to validation investigators
 - E. g. Overpass prediction
 - Try http://earthobservatory.nasa.gov/MissionControl/overpass.html